



## Clinical Policy: Double Balloon Enteroscopy

Reference Number: HNCA.CP.MP. 272

Effective Date: 5/06

Last Review Date: 3/19

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### Description

The double-balloon enteroscopy (DBE) is a deep small bowel endoscopic technique that allows for both diagnostic and therapeutic intervention of the entire small bowel, used primarily for suspected small bowel disease and bleeding.

### Policy/Criteria

- I. It is the policy of Health Net of California that DBE is **medically necessary** for the following indications:
  - A. Evaluation and treatment/therapeutic interventions for patients with obscure and/or occult gastrointestinal bleeding or suspected small bowel pathology, when esophagogastroduodenoscopy, colonoscopy and capsule endoscopy (CE) (or if CE is contraindicated) have failed to diagnose the source of bleeding.
  - B. A positive finding on CE requiring a biopsy or therapeutic intervention
  - C. For diagnosing suspected Crohn's disease when conventional diagnostic tests are negative
  - D. For removing entrapped foreign bodies in the small bowel (e.g., retained video capsule)
  - E. For tissue diagnosis and therapeutic interventions in patients with small-bowel tumors/malignancy or strictures detected by other diagnostic tests or in those with high suspicion for tumors despite initial negative testing.

### Background

The evaluation of the small bowel is difficult due to its length, intraperitoneal location, and contractility. A number of procedures used to assess this area include push enteroscopy, video capsule endoscopy, and intraoperative enteroscopy, which have advantages. However, limitations include decreased small bowel visualization, lack of therapeutic capacity, and a more invasive approach.

Bleeding from the small bowel is uncommon, but it is responsible for the majority of patients with gastrointestinal bleeding that persists or recurs without an obvious etiology after upper endoscopy, colonoscopy, and, possibly, radiologic evaluation of the small bowel. The evaluation of suspected small bowel bleeding is guided by the clinical history, physical findings, and the results of any previous evaluations. Additional tests that may be indicated include wireless video capsule endoscopy, deep small bowel enteroscopy, computed tomographic enterography or magnetic resonance enterography, and intraoperative enteroscopy. The most common first step in the evaluation of suspected small bowel bleeding is capsule endoscopy, provided the initial upper endoscopy and colonoscopy were completed with good visualization.<sup>1</sup>

DBE is an option to evaluate and treat small bowel disease or bleeding, and can be performed using either an antegrade or a retrograde approach. The insertion route is chosen according to



## CLINICAL POLICY

### Double Balloon Enteroscopy

the estimated location of the suspected lesions. This procedure is done when other conventional tests have failed to identify the source of bleeding or there is a contraindication to CE. <sup>5</sup>

There have been a number of studies, including a meta-analysis, as well as prospective, retrospective and case studies, which note that the diagnostic yield of DBE is equal or superior to that of push enteroscopy, fluoroscopic enteroclysis, small bowel radiography, and single-balloon enteroscopy. The evidence suggests that capsule endoscopy, if performed prior to DBE, increases the diagnostic yield. In addition, DBE can be used to treat certain small bowel lesions, particularly those responsible for occult bleeding, and thereby avoid other treatments, such as transfusion or surgery.

Kita et al (2018) in UptoDate notes that the most common indication for deep SBE is to evaluate suspected small bowel bleeding, and therapeutic indications for bleeding, polypectomy and stricture dilation. Serious complications can occur such as perforation and pancreatitis as with other endoscopy.

The American College of Gastroenterology guidelines (Syngal, et al., 2015), the European Society of Gastrointestinal Endoscopy (Pennazio, et al., 2015) and an Uptodate review (Chung and Adar, 2016) state that polyp removal thru device-assisted enteroscopy in patients with Peutz-Jeghers syndrome hasve been successfully performed but perforation risks may be increased with polypectomy due to serosal invagination within the polyp stalk.

#### *National Comprehensive Cancer Network*

There is no mention of DBE in these guidelines for either colon or gastric cancer.

#### *American College of Gastroenterology*

Video capsule endoscopy should be considered a first-line procedure for small bowel investigation, if there is no contraindication. Any method of deep enteroscopy can be used when endoscopic evaluation and therapy are required.

#### *Australian Medical Services Advisory Committee*

DBE is a safe, minimally invasive technique for examining endoscopically the whole of the small intestine, allowing biopsy and certain therapeutic procedures at the same time.

#### *American Society for Gastrointestinal Endoscopy (ASGE)(2015)*

In patients with suspected Crohn's disease, the overall yield of deep endoscopy (DE) for small-bowel pathology ranges from 30% to 48%, with an adverse event rate of approximately 1% for diagnostic examinations. A systematic review of diagnostic DBE for all indications found a pooled diagnostic yield of 63.4% (95% confidence interval, 42%-82.3%) for small-bowel pathology in patients with definite or suspected Crohn's disease, with a pooled minor and major adverse event rate for all indications of 9.1% and 0.72%, respectively.

DE is useful for tissue diagnosis and therapeutic interventions in patients with small-bowel tumors detected on video capsule endoscopy (VCE) or radiologic imaging. DE also is useful for detection of small-bowel tumors in patients in whom a high clinical suspicion for a tumor



## CLINICAL POLICY

### Double Balloon Enteroscopy

remains after negative VCE or other imaging. The miss rate of VCE for small-bowel tumors is reported as high as 18.9%, and malignant small-bowel tumors missed on VCE may be detected on DE. DE also may be useful for evaluating patients in whom VCE is contraindicated because of known or suspected small-bowel stenosis.

### Coding Implications

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2015, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

CPT® Codes	Description
44799	Unlisted procedure, small intestine

HCPCS Codes	Description
N/A	

### ICD-10-CM Diagnosis Codes that Support Coverage Criteria

ICD-10-CM Code	Description
C15.3-C15.8	Malignant neoplasm of esophagus
C17.0-C17.8	Malignant neoplasm of small intestine
C25.0-C25.8	Malignant neoplasm of pancreas
C49.A2	Gastrointestinal stromal tumor of stomach
C81.01-C81.19	Hodgkin lymphoma
C81.11-C81.19	Nodular sclerosis Hodgkin lymphoma
C81.21-C81.29	Mixed cellularity classical Hodgkin lymphoma
C81.31-C81.39	Lymphocyte depleted classical Hodgkin lymphoma
C81.41-C81.49	Lymphocyte-rich classical Hodgkin lymphoma
C81.71-C81.79	Other classical Hodgkin lymphoma
D50.0	Iron deficiency anemia secondary to blood loss (chronic)
D62	Acute posthemorrhagic anemia
D63.0	Anemia in neoplastic disease
K50.011-K50.018	Crohn's disease of small intestine, with complications (rectal bleeding, intestinal obstruction, fistula, abscess)
K57.00	Diverticulitis of small intestine with perforation and abscess without bleeding



## CLINICAL POLICY

### Double Balloon Enteroscopy

ICD-10-CM Code	Description
K57.01	Diverticulitis of small intestine with perforation and abscess, with bleeding
K57.10	Diverticulitis of small intestine with perforation and abscess
K57.11	Diverticulosis of small intestine without perforation or abscess, with bleeding
K57.12	Diverticulitis of small intestine without perforation or abscess, without bleeding
K57.13	Diverticulitis of small intestine without perforation or abscess with bleeding
K92.0	Hematemesis
K92.1	Melena
K92.2	Gastrointestinal hemorrhage, unspecified
T18.3X	Foreign body in the small intestine

Reviews, Revisions, and Approvals	Date	Approval Date
Policy Adopted from Health Net NMP#272 Double Balloon Enteroscopy	3/17	
Added indications C,D and E based on American Society for Gastrointestinal Endoscopy recommendations	3/18	3/18
Minor changes and references updated	3/19	3/19
No change, references updated	3/20	3/20

### References

1. Cave D. Evaluation of suspected small bowel bleeding (formerly obscure gastrointestinal bleeding). UpToDate. December 15, 2015.
2. Chauhan, SS et al. American Society for Gastrointestinal Endoscopy (ASGE) Technology Assessment Committee Technology Assessment Report on Enteroscopy. (2015) Accessed at: [https://www.asge.org/docs/default-source/education/Technology\\_Reviews/doc-enteroscopy.pdf](https://www.asge.org/docs/default-source/education/Technology_Reviews/doc-enteroscopy.pdf)
3. Chung DC, Adar T. Peutz-Jeghers syndrome: Screening and management. UpToDate Inc., Waltham, MA. Last reviewed July 2016.
4. Gerson L, Fidler JL, Cave DR, et al. American College of Gastroenterology. Clinical Guideline: Diagnosis and Management of Small Bowel Bleeding. Am J Gastroenterol 2015; 110:1265–1287. doi:10.1038/ajg.2015.246.
5. Hayes. Medical Technology Directory. Double-Balloon Video Enteroscopy for Diagnosis and Treatment of Small Bowel Disease. April 4, 2014. Archived May 28, 2015.
6. Kaffes AJ, Koo JH, Meredith C, et al. Double-balloon enteroscopy in the diagnosis and the management of small-bowel diseases: an initial experience in 40 patients. Gastrointest Endosc. 2006; 63(1):81.
7. Khashab, MA et al American Society for Gastrointestinal Endoscopy (ASGE) Standards of Practice Committee The role of deep enteroscopy in the management of small bowel disorders. (2015) Accessed at: <https://www.asge.org/docs/default->



## CLINICAL POLICY

### Double Balloon Enteroscopy

[source/education/practice\\_guidelines/doc-deep\\_enteroscopy\\_small\\_bowel\\_disorders\\_aip.pdf?sfvrsn=6](#)

March 2018

8. Kita H. Overview of deep small bowel enteroscopy. UpToDate. Updated June 27, 2016 and June 2018,
9. Medical Services Advisory Committee (MSAC). Double balloon enteroscopy. Assessment Report. MSAC Application No. 1102. Canberra, ACT: MSAC; 2006. Available at: <http://www.msac.gov.au/internet/msac/publishing.nsf/Content/app1102-1>.
10. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Gastric Cancer. Version 3.2016.
11. National Comprehensive Cancer Network (NCCN). NCCN Clinical Practice Guidelines in Oncology. Colon Cancer. Version 1.2017.
12. Pasha SF, Leighton JA, Das A, et al. Double-balloon enteroscopy and capsule endoscopy have comparable diagnostic yield in small-bowel disease: a meta-analysis. *Clin Gastroenterol Hepatol*. 2008 Jun;6(6):671-6. doi: 10.1016/j.cgh.2008.01.005. Epub 2008 Mar 20.
13. Pennazio M, Spada C, Eliakim R, et al. Small-bowel capsule endoscopy and device-assisted enteroscopy for diagnosis and treatment of small-bowel disorders: European Society of Gastrointestinal Endoscopy (ESGE) Clinical Guideline. *Endoscopy*. 2015;47(4):352-376.
14. Rahman A, Ross A, Leighton JA, et al. Double-balloon enteroscopy in Crohn's disease: findings and impact on management in a multicenter retrospective study. *Gastrointest Endosc*. 2015 Jul;82 (1):102-7. doi: 10.1016/
15. Robles EP, Delgado PE, Conesa PB, et al. Role of double-balloon enteroscopy in malignant small bowel tumors. *World J Gastrointest Endosc*. 2015; 7(6):652-658.
16. Samaha E, Rahmi G, Landi B, et al. Long-term outcome of patients treated with double balloon enteroscopy for small bowel vascular lesions. *Am J Gastroenterol*. 2012 Feb; 107(2):240-6. doi: 10.1038/ajg.2011.325. Epub 2011 Sep 27.
17. Sun B, Rajan E, Cheng S, et al. Diagnostic yield and therapeutic impact of double-balloon enteroscopy in a large cohort of patients with obscure gastrointestinal bleeding. *Am J Gastroenterol*. 2006; 101(9):2011.
18. Teshima CW, Kuipers EJ, van Zanten SV, et al. Double balloon enteroscopy and capsule endoscopy for obscure gastrointestinal bleeding: An updated meta-analysis. *J Gastroenterol Hepatol*. 2011;26(5):796-801.
19. Travis AC, Saltzman JR. Evaluation of Occult Gastrointestinal Bleeding. UpToDate. Oct.24, 2016.
20. Wadwa V, Sethi S, Tewani S, et al. A meta-analysis on efficacy and safety: single-balloon vs. double-balloon enteroscopy. *Gastroenterol Rep*. 2015 May;3(2):148-55. doi: 10.1093/gastro/gov003. Epub 2015 Feb 18.

### **Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and



## CLINICAL POLICY

### Double Balloon Enteroscopy

accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

**Note: For Medicaid members**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.



Health Net®

## CLINICAL POLICY

### Double Balloon Enteroscopy

**Note: For Medicare members,** to ensure consistency with the Medicare National Coverage Determinations (NCD) and Local Coverage Determinations (LCD), all applicable NCDs and LCDs should be reviewed prior to applying the criteria set forth in this clinical policy. Refer to the CMS website at <http://www.cms.gov> for additional information.

©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.